

ARTICLE

Received 6 Apr 2010 | Accepted 17 Nov 2010 | Published 14 Dec 2010

DOI: 10.1038/ncomms1139

High rates of photobiological H₂ production by a cyanobacterium under aerobic conditions

Anindita Bandyopadhyay¹, Jana Stöckel¹, Hongtao Min², Louis A. Sherman² & Himadri B. Pakrasi¹

Among the emerging renewable and green energy sources, biohydrogen stands out as an appealing choice. Hydrogen can be produced by certain groups of microorganisms that possess functional nitrogenase and/or bidirectional hydrogenases. In particular, the potential of photobiological hydrogen production by oxygenic photosynthetic microbes has attracted significant interest. However, nitrogenase and hydrogenase are generally oxygen sensitive, and require protective mechanisms to function in an aerobic extracellular environment. Here, we describe *Cyanothece* sp. ATCC 51142, a unicellular, diazotrophic cyanobacterium with the capacity to generate high levels of hydrogen under aerobic conditions. Wild-type *Cyanothece* 51142 can produce hydrogen at rates as high as 465 μmol per mg of chlorophyll per hour in the presence of glycerol. Hydrogen production in this strain is mediated by an efficient nitrogenase system, which can be manipulated to convert solar energy into hydrogen at rates that are several fold higher, compared with any previously described wild-type hydrogen-producing photosynthetic microbe.

¹ Department of Biology, Washington University, St Louis, Missouri 63130, USA. ² Division of Biological Sciences, Purdue University, W. Lafayette, Indiana 47907, USA. Correspondence and requests for materials should be addressed to H.B.P. (email: pakrasi@wustl.edu).

Microbial H₂ production relies on either photosynthetic or fermentative processes. Economic feasibility studies suggest that a direct and efficient conversion of solar energy to H₂ in a carbon-neutral way is necessary for the process to be commercially viable^{1,2}. Several photosynthetic microalgal and bacterial species possessing nitrogenase and/or hydrogenase enzymes are being studied as prospective model organisms for photobiological H₂ production^{3–5}. Outstanding among them are *Rhodospseudomonas palustris*, a purple photosynthetic, nitrogen-fixing bacterium; *Chlamydomonas reinhardtii*, a green eukaryotic alga; as well as *Anabaena* and *Synechocystis*, members of the cyanobacterial group. High specific rates of nitrogenase-mediated H₂ production have been reported for *R. palustris*⁶. However, *R. palustris* performs anoxygenic photosynthesis, thereby requiring an anaerobic environment for photobiological H₂ production^{7–9}. *Anabaena*, a filamentous diazotrophic cyanobacterial strain, produces H₂ in heterocysts, specialized N₂-fixing cells, which maintain a microaerobic environment to protect the oxygen-sensitive nitrogenase enzyme. However, the low frequency of heterocysts in a filament (about one in ten cells normally differentiates into heterocysts¹⁰) consequently results in modest yields of net H₂ production. In contrast, H₂ production in *C. reinhardtii* and *Synechocystis* sp. PCC 6803 is mediated by hydrogenase enzymes, and H₂ production can be achieved in both of these organisms only under strictly anaerobic conditions^{11–13}. Although these strains have long been used as model organisms to study biohydrogen production, the importance of selecting additional novel and native strains with diverse energy conversion systems that might have evolved as a consequence of specific ecological pressure has often been emphasized^{1,14}. A recent effort in this direction has identified ten native N₂-fixing, heterocystous cyanobacterial strains that exhibit higher rates of H₂ production compared with some of the previously studied mutant strains¹⁵.

In the present report, we describe *Cyanothece* sp. ATCC 51142, a cyanobacterial strain with the ability to produce remarkably high amounts of H₂ under aerobic conditions. *Cyanothece* strains thrive in marine environments limited in dissolved inorganic bioavailable nitrogenous compounds, and have been recognized for their role in maintaining the marine nitrogen cycle^{16–18}. *Cyanothece* 51142 can derive the majority of its nutritional requirements from sunlight, atmospheric carbon dioxide and nitrogen gases (Fig. 1). During the day, it performs photosynthesis and fixes carbon, which is stored as large reserves of glycogen¹⁹. At the onset of the dark period, high rates of respiration rapidly create a suboxic intracellular environment. This, in turn, facilitates oxygen-sensitive and energy-intensive processes such as N₂ fixation and H₂ production to occur at night at the expense of the accumulated glycogen. The orchestrated diurnal cycling patterns of the central metabolic processes in this organism have recently been described using a global transcriptomic approach²⁰. These unique attributes of *Cyanothece* 51142, which make it an ideal organism for H₂ production, are possibly the remnants of the metabolic and regulatory processes that aided in the acclimatization of ancient cyanobacteria during their transition from an anaerobic to an aerobic environment. Retention of ancient metabolic traits that originated in the Archaean oceans has been reported in other cyanobacterial strains²¹.

Results

A two-stage system for photobiological H₂ production. We developed a two-stage H₂ production system in *Cyanothece* 51142 on the basis of our previous knowledge of the diurnal rhythms in this unicellular cyanobacterium^{22,23}. The two stages were comprised of a growth phase during which cells were allowed to grow aerobically under 12 h light/12 h dark cycles, followed by an incubation phase during which cells sampled at the end of the 12 h light period were incubated in airtight vials under continuous illumination for 12 h. During this latter light-incubation period, the physiological activities




Figure 1 | Photobiological H₂ production in *Cyanothece* 51142. A schematic diagram showing the process of biohydrogen production by *Cyanothece* 51142 cells using solar energy and atmospheric CO₂ and/or glycerol. CO₂ is fixed during the day to synthesize glycogen, which serves as an energy reserve and electron source for H₂ production at night.

of the cells were in step with the subjective dark condition, thereby facilitating N₂ fixation and H₂ production. At the completion of the incubation phase, the headspace of the vials (which contained 100% air at the beginning of incubation) was analysed for H₂ accumulation. We determined that photoautotrophically grown *Cyanothece* 51142 exhibited high specific rates of H₂ production (> 150 μmol of H₂ per mg of chlorophyll per hour (Chl.h) or 2.34 μmol of H₂ per mg of dry weight per hour) under aerobic incubation conditions (Fig. 2a). This is striking, as most unicellular photosynthetic microbial strains require a complete anaerobic environment for H₂ production²⁴ (Table 1).

The rates of H₂ production in *Cyanothece* 51142 could be greatly enhanced when cells were grown in the presence of additional carbon sources, as observed in cultures supplemented with high concentrations of CO₂, or glycerol (Fig. 2a). Cells grown under CO₂-enriched air and incubated under aerobic conditions could produce 230 μmol of H₂ per mg of Chl.h (Fig. 2a). In fact, a batch culture of *Cyanothece* 51142 cells supplemented with 50 mM glycerol could produce more than 900 ml of H₂ per litre of culture over a period of 2 days (Fig. 2b,c) when incubated in constant light under aerobic conditions (Fig. 2b). Notably, the kinetics of H₂ production under these conditions revealed high rates even early in the incubation phase when the O₂ concentration in the headspace of the vials was equivalent to that in air (Fig. 2b), indicating that an intracellular anoxic environment facilitates nitrogenase activity during the subjective dark period. The O₂ level in the headspace diminished with time (Fig. 2b), indicating continued respiratory activities in the cells. Moreover, as shown in the next subsection, transcripts of *coxA*, the gene encoding subunit A of the cytochrome *c* oxidase enzyme involved in respiration, were also detected under these conditions. This respiration-induced microaerobic environment was also reflected in the dissolved O₂ levels in the culture in the incubation vials (Fig. 2d). A rapid decline in the dissolved O₂ concentration was observed at the beginning of the incubation phase, followed by periodic oscillations in O₂ levels, which corresponded to the respiratory and photosynthetic activity of the cells in the subjective dark and light periods, respectively. In comparison with light incubation,




Figure 2 | Aerobic H₂ production rates and generation of a suboxic intracellular environment in *Cyanosethece* 51142. (a) Specific rates of H₂ production by *Cyanosethece* 51142 grown under aerobic, photoautotrophic conditions using ambient (0.03%, v/v) or high (8%, v/v) concentrations of CO₂, and under photomixotrophic growth conditions with glycerol (50 mM). Cultures were incubated under aerobic conditions for H₂ production. Each column represents an average of measurements from at least three biological replicates. Error bars indicate s.d. values from the average. (b) Kinetics of aerobic photobiological H₂ production in a batch culture of *Cyanosethece* 51142. H₂ production (blue) could be observed for more than 60 h in vials containing cultures under aerobic incubation conditions in the presence of light. The respiratory activities of the cells resulted in a gradual decrease in O₂ concentration in the headspace of the vials (red). Three biological replicates were used for the measurements. Error bars indicate s.d. values from the average. (c) Comparison of cumulative H₂ accumulation in vials with cells under light and dark incubation conditions. A volume of 916 ml of H₂ per litre of culture accumulated in the vials when incubated in the light compared with 7 ml of H₂ per litre of culture accumulation in the dark. Each column represents an average of measurements from at least three biological replicates. Error bars indicate s.d. values from the average. (d) Dissolved O₂ levels in a batch culture of *Cyanosethece* 51142 cells under H₂-producing conditions. The circadian rhythms of respiratory and photosynthetic activities were reflected in the dissolved O₂ concentration of the culture. The culture was transferred to incubation vials at the beginning of the dark period when the dissolved O₂ level was about 50% of the maximum (assuming 100% equals 230 μM dissolved O₂ at 30 °C). The O₂ concentration reduced starting from the middle of the subjective light period (blue line, grey bar), was at its minimum at the middle of the subjective dark period (red line, white bar), after which an increase was observed because of photosynthetic oxygen evolution. The horizontal bars below the x axis denote subjective day (open) and night (filled) periods. Three independent biological replicates were used for the measurements. Error bars indicate s.d. values from the average.

incubation under dark conditions reduced the level of H₂ production by more than 100-fold, indicating that H₂ production in this system is driven by light energy, and the high rates observed cannot be achieved by dark fermentation of glycerol alone (Fig. 2c).

Additional carbon sources lead to higher glycogen content. We assessed the influence of external carbon sources on the cellular energy reserves, a critical determinant of nitrogenase activity and H₂ production. The glycogen levels of the cells were compared between

pre- (I₀) and post-incubation (I₁₂) samples collected from cultures grown under ambient CO₂ (0.03%), under 8% CO₂-enriched air, and with 50 mM glycerol. Either high CO₂ or glycerol enhanced the glycogen reserves of the cells (Fig. 3). In ambient and high CO₂-treated cells, the glycogen level diminished at the end of the incubation period, indicating that the endogenous carbon source was being used for H₂ production (Fig. 2a). In contrast, cells grown in the presence of glycerol showed higher levels of glycogen at the end of the 12 h-light incubation phase (I₁₂), implying direct utilization of glycerol as a source of ATP and reductants by the nitrogenase enzyme under this photomixotrophic condition. This is supported by the observation that expression of glycogen phosphorylase (*glgP*), a gene involved in glycogen degradation, was also down-regulated in glycerol-supplemented cells (Fig. 4b). In this context, we have recently shown that *Cyanosethece* 51142 uses glycerol as the sole carbon source when cells are grown under photomixotrophic conditions in the presence of glycerol²⁵.

Hydrogen production is mediated by the nitrogenase enzyme.

Genome analysis of *Cyanosethece* 51142 revealed the presence of both the nitrogenase (Nif) and bidirectional hydrogenase (Hox) enzyme systems. Analysis at the transcriptional level revealed that *hox* genes were expressed under both nitrogen-fixing (–NO₃) and nitrogen-sufficient (+NO₃) conditions (Fig. 4a), and higher transcript abundance could be associated with the dark period under both conditions. In contrast, *nif* genes were expressed only under nitrogen-fixing conditions, and, in the absence of any additional carbon source, the *nif* transcripts were strictly associated with the dark period (Fig. 4a). High levels of H₂ production in *Cyanosethece* 51142 were detected exclusively under diazotrophic conditions, and addition of combined nitrogen to the growth media of nitrogen-fixing cultures resulted in an immediate reduction in the rates of nitrogen fixation and simultaneous cessation of H₂ production (Fig. 4c). These results show that the high rates of H₂ production observed in *Cyanosethece* 51142 were primarily mediated by the nitrogenase enzyme system.

Nitrogenase can be optimized to enhance hydrogen yield. An important feature of the nitrogenase enzyme system is the potential to channel all available electrons towards H₂ production in the absence of molecular nitrogen^{26,27}. The nitrogenase reaction is also resistant to feedback inhibition from accumulated H₂ (ref. 28). When *Cyanosethece* 51142 cells grown under photoautotrophic conditions were incubated under an argon atmosphere (absence of molecular nitrogen), a 2- to 3-fold increase (> 370 μmol of H₂ per mg of Chl.h or 5.738 μmol of H₂ per mg of dry weight per hour) in the yield of H₂ was achieved (Fig. 5, Tables 1 and 2). In addition, production rates of up to 467 μmol of H₂ per mg of Chl.h were achieved from cells grown in glycerol-supplemented media subsequently incubated under argon environment (Fig. 5). These rates are at least an order of magnitude higher compared with any other hydrogen-producing photosynthetic microbial wild-type strain studied to date (Tables 1 and 2).

Discussion

This work describes the use of the wild-type strain of the unicellular cyanobacterium *Cyanosethece* 51142 for highly efficient photobiological H₂ production under natural aerobic conditions. In general, the oxygen sensitivity of the enzymes involved in biological H₂ production makes the use of oxygenic photosynthetic organisms as a platform for H₂ production an extremely challenging task. Thus, until now, photobiological H₂ production studies have largely relied on artificial interventions, which help to create an anaerobic environment. It has been recognized that the use of oxygenic photosynthetic microbes that can produce H₂ under aerobic conditions would be an important step forward for biological H₂ production²⁹.

Table 1 | Specific rates of H₂ production by wild-type strains of *Cyanothece* 51142 and other model photosynthetic microbes.

Strain	Enzyme	Specific rates (aerobic incubation)		Specific rates (anaerobic incubation)	
		($\mu\text{mol per mg of Chl.h}$)	($\mu\text{mol per mg of Chl.h}$)	($\mu\text{mol per mg of Chl.h}$)	($\mu\text{mol per mg of protein h}^{-1}$)
<i>Cyanothece</i> 51142	Nitrogenase	152	373	3.5	
<i>Anabaena</i> 29413	Nitrogenase	—	39.4 (ref. 24)	0.02 (ref. 42)	
<i>Synechocystis</i> 6803	Hydrogenase	—	1.2* (ref. 11)	ND	
<i>Chlamydomonas reinhardtii</i>	Hydrogenase	—	6.6*† (refs 12, 13)	ND	
<i>Rhodospseudomonas palustris</i>	Nitrogenase	ND	ND	0.92† (refs 9, 28)	

Abbreviations: ND, not determined; —, not detected.
 *Initial rates of H₂ production.
 †The rates were calculated using information from references 9, 28, 12 and 13.




Figure 3 | Effects of external carbon sources on glycogen accumulation in *Cyanothece* 51142. I₀ (red) and I₁₂ (blue) indicate the beginning and end of a 12 h-light incubation period for H₂ production, respectively. Differences in glycogen level between the two time points correspond to the amount of glycogen used for N₂ fixation/H₂ production. Each column represents an average of measurements from at least three biological replicates. Error bars indicate s.d. values from the average.

Our study has identified *Cyanothece* 51142 as one such organism that has developed an effective strategy to make the best use of a diurnal cycle—synthesizing energy-rich storage compounds during the day and using it for nitrogen fixation at night when oxygen-consuming processes render the interior of the cell anaerobic (or sub-oxic), while the extracellular environment continues to be oxygen rich. This trait of *Cyanothece* 51142 formed the basis for the physiological perturbations that were designed for our two-stage aerobic H₂ production process.

Nitrogen fixation is an energy-intensive process, requiring 16 molecules of ATP for every molecule of nitrogen fixed and H₂ produced¹⁹ (Fig. 1). However, the process is of paramount importance to diazotrophic species inhabiting ecological niches (such as deep oceans) with very low levels of nitrogenous nutrients. Consequently, photoautotrophic unicellular strains such as *Cyanothece* 51142 are expected to have evolved effective strategies for collecting and storing solar energy, which can be used at night when the energy demands are high. Our study shows that *Cyanothece* 51142 not only develops an intracellular environment conducive for the function of the nitrogenase enzyme but also generates an adequate supply of ATP for this high-energy-requiring process. In addition, our results reveal high specific activity of the nitrogenase enzyme in this strain, a finding consistent with earlier reports that showed higher rates of nitrogen fixation in marine unicellular diazotrophs such as *Cyanothece* compared with some filamentous strains^{16,30}. A unicellular, marine, diazotrophic *Synechococcus* strain (*Synechococcus* Miami

BG 43511; later classified as *Cyanothece* Miami BG 43511¹⁶) was also shown to exhibit high rates of nitrogenase activity and hydrogen production³¹. Unfortunately, this strain has not been readily accessible to the general research community. Notably, not all unicellular cyanobacterial strains are endowed with these traits, as is evident from studies on *Gloeotheca*, a unicellular freshwater diazotroph that possesses a nitrogenase enzyme with relatively low specific activity and does not have any appreciable nitrogenase-mediated H₂ production capacity³².

Diazotrophic cyanobacteria have developed various strategies to protect their nitrogenase enzyme from the oxygen-rich environment they inhabit. However, unlike *Cyanothece* 51142, most diazotrophic strains are unable to exhibit nitrogenase-mediated H₂ production under aerobic conditions. In filamentous, heterocyst-forming diazotrophic strains, this is largely ascribed to the activities of an uptake hydrogenase enzyme system that is functionally closely associated with nitrogenase and oxidizes the H₂ produced^{3,33–35}. It has been shown that wild-type *Anabaena variabilis* cells can generate H₂ only under an argon atmosphere, whereas its uptake hydrogenase mutants PK84 and AVM13 can produce H₂ aerobically^{27,36–38}. A recent study also demonstrated H₂ production (~25 $\mu\text{mol per mg of Chl.h}$) from the vegetative cells of wild-type *Anabaena variabilis* under nitrogen atmosphere when strict anaerobic conditions are maintained³⁹. The genome sequence of *Cyanothece* 51142 shows the presence of *hup* genes for an uptake hydrogenase. The transcripts for one of these genes, *hupS*, were also detected under H₂-producing conditions (Fig. 4b). Interestingly, the *hupS* transcripts in *Cyanothece* 51142 were present under both nitrogen-sufficient and nitrogen-fixing conditions (Fig. 4a), indicating that its expression is independent of *nif*. The presence of *hupS* transcripts and the concurrent accumulation of H₂ at high rates under aerobic incubation conditions are suggestive of a weak uptake hydrogenase activity in *Cyanothece* 51142. Such a premise is also supported by the observation that a few wild-type *Anabaena* strains that possess uptake hydrogenases with very low specific activities can also exhibit aerobic H₂ production^{24,40,41}.

The ability to use high concentrations of CO₂ or glycerol for enhanced H₂ production is an added advantage, as both of these carbon sources are abundantly available as industrial waste products, making biohydrogen production by *Cyanothece* 51142 an attractive option. High CO₂ and glycerol provide an additional carbon source and the availability of excess carbon functions as a signal for enhanced nitrogenase activity to meet the increase in nitrogen demand in these cyanobacterial cells. The rise in the glycogen level of cells at the end of the incubation phase in glycerol-supplemented cultures could be a result of cellular activities geared towards building energy reservoirs when an external energy source is readily available²¹.

Decades of research have unveiled various principles underlying biological H₂ production. However, achieving significant increases in yield has been a major challenge. Genetic modifications of H₂-yielding pathways have resulted in improvements in production




Figure 4 | H₂ production in *Cyanothecce* 51142 is primarily mediated by the nitrogenase enzyme system. (a) Comparisons of the expression of genes involved in H₂ metabolism under N₂-fixing (-NO₃) and nitrogen-sufficient (+NO₃) conditions from cultures grown under 12h light/dark cycles. *nifH* transcripts could only be observed under N₂-fixing condition, whereas *hoxH* and *hupS* transcripts were present under both conditions. *coxA* transcripts were predominant during the dark cycle both under N₂-sufficient and N₂-deficient conditions. Transcripts for *glgP* were abundant during the dark cycle under both conditions. *16S rRNA* was used as the loading control. Cells were collected at 4h intervals, between D1 (1h after the onset of the dark cycle) and L9 (9h into the light cycle). (b) Reverse transcription PCR analysis of the enzyme systems involved in H₂ metabolism in *Cyanothecce* 51142. Samples were collected from the incubation bottles (continuous light, aerobic) at the beginning (I₀) and end of the incubation phase (I₁₂). Strong *nifH* expression could be observed at I₀ in the control (grown in ambient CO₂) as well as in cultures supplemented with high concentrations of CO₂ or glycerol. *nifH* transcripts were also present in cells incubated under light during H₂ production (I₁₂). Expression of *glgP* was downregulated in cultures supplemented with glycerol, indicating that glycerol, and not glycogen, is used directly as a carbon source under these mixotrophic conditions. Interestingly, *hoxH* expression was also significantly downregulated in the presence of glycerol. Expression of *hupS* was lower under light. The presence of *coxA* transcripts suggested respiratory activities in the vials under all conditions. *16S rRNA* was used as the loading control. -RTase denotes controls to demonstrate absence of DNA contamination in the samples. (c) Addition of NaNO₃ to the medium of an N₂-fixing culture resulted in cessation of H₂ production (blue) within 12h and greatly reduced nitrogen fixation (green). Nitrogenase activity is expressed in terms of ethylene production by acetylene reduction. Each column represents an average of measurements from at least three biological replicates. Error bars indicate s.d. values from the average.




Figure 5 | Effect of argon environment on H₂ production in *Cyanothecce* 51142. Higher specific rates of H₂ production are achieved in the absence of molecular N₂ in the headspace of the incubation bottles. Each column represents an average of measurements from at least three biological replicates in air (blue) and argon (red) atmospheres. Error bars indicate s.d. values from the average.

rates compared with the corresponding wild-type strains^{9,42,43}. However, as the H₂ production rates in these wild-type strains are rather modest, even a 20-fold increase in yield in mutant strains is not sufficient to attain a high production level. Therefore, our identification of a cyanobacterial strain exhibiting high rates of H₂ production under ambient aerobic conditions offers new possibilities in photobiological hydrogen production research. Recent studies have revealed the metabolic flexibility of this cyanobacterium²⁵, and demonstrated that its robust circadian rhythm allows N₂ fixation

and H₂ production to occur at reasonably high rates even when grown under continuous light⁴⁴. Previous studies have shown the robustness of other cyanobacterial systems for H₂ production over a prolonged period of time³⁸, demonstrating the possibility of using high-H₂-yielding cyanobacterial strains for large-scale production. A systems level understanding of this biological phenomenon in *Cyanothecce* 51142 will unravel previously unknown cellular factors and regulatory mechanisms that influence the process so that they can be favourably altered to produce even higher levels of H₂ as an energy carrier.

Methods

Growth conditions. For H₂ measurement, *Cyanothecce* 51142 cells were grown in shaking flasks in ASP2 medium⁴⁵ without supplemented NaNO₃ at 30°C under 12h light/12h dark cycles and 100 μmol photons per m² s⁻¹ of white light. Cultures were inoculated with 0.25 volumes of cultures grown in ASP2 medium without NaNO₃ under continuous light (50 μmol photon per m² s⁻¹ white light), which in turn were inoculated with 0.1 volumes of cultures grown in ASP2 with NaNO₃ under similar conditions. For photomixotrophic growth, cultures were supplemented with 50 mM glycerol. For growth under high-CO₂ conditions, the cultures were aerated with 8% CO₂-enriched air at a flow rate of 100 ml min⁻¹.

H₂ production and nitrogenase activity measurement. A volume of 20 ml of culture was transferred at the beginning of the dark period to air-tight glass vials (36 ml) and incubated in air under a light intensity of 100 μmol photon per m² s⁻¹ for 12h. The chlorophyll content of cultures grown without an external carbon source ranged between 0.5 and 2 μg ml⁻¹, whereas cultures supplemented with glycerol had higher chlorophyll concentration (2–5 μg ml⁻¹). Dry weight of photoautotrophically grown culture ranged between 90 and 98 μg ml⁻¹. For anaerobic incubation, the glass vials were flushed with argon for 15–30 min. For the batch culture experiment, 25 ml of dense (8–12 μg chlorophyll per ml culture) glycerol-supplemented culture was incubated in 145 ml vials under constant illumination. H₂ that accumulated in the headspace of sealed culture vials was withdrawn with

Table 2 | Improvements in the rates of H₂ production by physiological and genetic modifications.

Strain	Specific rates of H ₂ production		Conditions
	($\mu\text{mol per mg of protein h}^{-1}$)	($\mu\text{mol per mg of Chl.h}$)	
<i>Cyanothece</i> 51142	3.5	373	WT, photoautotrophic growth under ambient CO ₂ concentrations, argon incubation
<i>Anabaena</i> 29413 (PK84 mutant)	0.62 (ref. 42)	167.6 (ref. 24)	<i>hup</i> ⁻ mutant ¹ , 2% CO ₂ , argon incubation ²⁴
<i>Synechocystis</i> 6803 (M55 mutant)	ND	5* 6†	<i>ndhB</i> ⁻ mutant ¹ , glucose, glucose oxidase, sulphur deprivation, argon incubation ¹¹
<i>Chlamydomonas reinhardtii</i>	ND	16.26*‡ 5.8	WT, photomixotrophic growth, sulphur deprivation, anaerobic incubation ^{12,13}
<i>Rhodospseudomonas palustris</i>	3.6	ND	Mutations in <i>hup</i> and <i>nifA</i> , organic carbon sources, anaerobic growth and anaerobic incubation ²⁸

Abbreviations: ND, not determined; WT, wild-type.
 *Initial rates of H₂ production.
 †Rates last for > 25 min.
 ‡The rates were calculated using information from references 12 and 13.
 ||Rates last for > 55 h.

an air-tight syringe and quantified using an Agilent 6890N Gas Chromatograph (Agilent) equipped with a Molsieve 5A 60/80 column (Molsieve; inner dimensions 6'×1/8") and a thermal conductivity detector. Injection port, oven and detector temperatures were 100, 50 and 100 °C, respectively. Argon, the carrier gas, was supplied at a flow rate of 65 ml min⁻¹. The volume of gas expressed in the results section was under standard conditions (assuming 1 ml H₂ = 44.6 μmol).

Nitrogenase activity of the H₂-producing cultures was determined using an acetylene reduction assay⁴⁶ and expressed in terms of the ethylene produced. Cells were incubated in sealed glass vials in light at 30 °C under a 5% acetylene atmosphere with or without flushing with argon. Gas samples were withdrawn, and ethylene production was measured using an Agilent 6890N Gas Chromatograph (Agilent) equipped with a Poropak N column (inner dimensions 5'×1/8") and a flame ionization detector using argon as the carrier gas (flow rate of 65 ml min⁻¹), according to the manufacturer's instructions. The temperature of the injector, detector and oven were 150, 200 and 100 °C, respectively.

Total chlorophyll *a* was extracted by methanol and quantified spectrophotometrically using an Olis DW2000 spectrophotometer (On-Line Instrument Systems). Protein concentrations were determined using a bicinchoninic acid assay (Pierce) according to the manufacturer's instructions.

Oxygen measurements. O₂ concentrations in the headspace of the incubation vials were determined using an Agilent 6890N Gas Chromatograph (Agilent) equipped with a Molsieve 5A 60/80 column (Molsieve; inner dimensions 6'×1/8") and a thermal conductivity detector using the same settings as described above for the H₂ assay. The measurements were taken under standard conditions. Dissolved O₂ concentration in the incubation vials was measured using a Clark-type electrode. Calculations were based on the fact that air-saturated water contains 230 μM of O₂ at 30 °C.

Determination of glycogen content. Samples were collected for the glycogen assay at the beginning and end of the 12 h-light incubation phase of H₂ production. The cellular glycogen content was measured using a glucose hexokinase assay (Sigma) with glycogen from bovine liver Type IX (Sigma) as standard. After methanol extraction of chlorophyll, the cell pellets were washed twice with 100% ethanol. To remove free glucose, 40% KOH was added and the samples were incubated for 1 h at 95 °C. Glycogen was precipitated overnight at -20 °C with 2 volumes of 100% ethanol. The samples were centrifuged for 1 h at 4 °C and 2 N HCl was added before incubation at 95 °C for 30 min. The same volume of 2 N NaOH and 0.5 volumes of 1 M phosphate buffer, pH 7, were added before dilution with 1 volume of distilled water. For the enzyme assay, 75 μl of sample solution was mixed with 200 μl of enzyme solution in a microtitre plate (Costar, ultraviolet light proof). After 15 min incubation at ambient temperature, NADPH was measured at 340 nm on a μQuant plate reader (Bio-Tek Instruments).

Semiquantitative reverse transcription (RT-PCR). Semiquantitative RT-PCR analyses were performed on RNA samples isolated from cultures grown under nitrogen-fixing (-NO₂) conditions, with and without supplemented glycerol; from cultures grown under aeration with 8% CO₂-enriched air; and under non-nitrogen-fixing conditions (+NO₂). For the time-course experiment, samples were collected every 4 h for 24 h, starting with 1 h into the dark period (D1). In total, six samples were collected. For RT-PCR analysis under H₂-producing conditions, culture samples were assayed at the end of the light period at time point I₀ and from the assay

bottles at the end of incubation at time point I₂. RNA was isolated and quantified essentially as described in ref. 14. A volume of 700 ng of DNase (Promega)-treated total RNA samples was used for reverse transcription with the Superscript II Reverse Transcriptase and random primers (Invitrogen) according to the manufacturer's instructions. The absence of DNA contamination was tested for each RNA sample (Fig. 4b). PCR was carried out at 94 °C for 4 min, followed by 94 °C for 30 s, 58 °C for 20 s, 72 °C for 20 s and a final extension time of 4 min at 72 °C. A total of 25 cycles for the *nifH*, *hupS*, *glgP* and *coxA* genes and 26 cycles for the *hoxH* gene were used. The following primers were used: *nifH* F: 5'-ACCATT GCTGCGTTAGCTGAAAC-3', R: 5'-TAATACCACGACCCGCACATCCA-3'; *coxA* F: 5'-TGATATGGCCTTTCCACCCTCA-3', R: 5'-AGAGAAGCTAAAGCG GCAGCGAGA-3'; *hupS* F: 5'-ATAGCTGGTTTCGTTGTCGCTGT-3', R: 5'-CGAAGTCTTGGGTGGTTGCTTTG-3'; *hoxH* F: 5'-TGGAGAAGACG GACTTTGGGAAC-3', R: 5'-AAAGAAGAGGTGCTACACCACC-3'; *glgP* F: 5'-TCGGCTGAATTCCTTATGGGTCG-3', R: 5'-CAGGAATTTCCACTT GCCAACCC-3'; *16S rRNA* F: 5'-AGAGGATGAGCAGCCACT-3', R: 5'-TAAT TCCGGATAACGCTTGC-3' (F: forward, R: reverse).

References

- Carrieri, D., Kolling, D., Ananyev, G. & Dismukes, G. C. Prospecting for biohydrogen fuel. *Indst. Biotechnol.* **2**, 133–137 (2006).
- Das, D. & Veziroglu, T. N. Advances in biological hydrogen production processes. *Int. J. Hydrogen Energy* **33**, 6046–6057 (2008).
- Tamagnini, P. *et al.* Hydrogenases and hydrogen metabolism of cyanobacteria. *Microbiol. Mol. Biol. Rev.* **66**, 1–20 (2002).
- Ghirardi, M. L. *et al.* Hydrogenases and hydrogen photoproduction in oxygenic photosynthetic organisms. *Annu. Rev. Plant Biol.* **58**, 71–91 (2007).
- Tamagnini, P. *et al.* Cyanobacterial hydrogenases: diversity, regulation and applications. *FEMS Microbiol. Rev.* **31**, 692–720 (2007).
- Melnicki, M. R., Bianchi, L., De Philippis, R. & Melis, A. Hydrogen production during stationary phase in purple photosynthetic bacteria. *Int. J. Hydrogen Energy* **33**, 6525–6534 (2008).
- Barbosa, M. J., Rocha, J. M. S., Tramper, J. & Wijffels, R. H. Acetate as a carbon source for hydrogen production by photosynthetic bacteria. *J. Biotechnol.* **85**, 25–33 (2001).
- Gosse, J. L. *et al.* Hydrogen production by photoreactive nanoporous latex coatings of nongrowing *Rhodospseudomonas palustris* CGA009. *Biotechnol. Prog.* **23**, 124–130 (2007).
- Rey, F. E., Heiniger, E. K. & Harwood, C. S. Redirection of metabolism for biological hydrogen production. *Appl. Environ. Microbiol.* **73**, 1665–1671 (2007).
- Golden, J. W. & Yoon, H. S. Heterocyst formation in *Anabaena*. *Curr. Opin. Microbiol.* **1**, 623–629 (1998).
- Cournac, L., Guedeney, G., Peltier, G. & Vignais, P. M. Sustained photoevolution of molecular hydrogen in a mutant of *Synechocystis* sp strain PCC 6803 deficient in the type I NADPH-dehydrogenase complex. *J. Bacteriol.* **186**, 1737–1746 (2004).
- Kosourov, S., Tsygankov, A., Seibert, M. & Ghirardi, M. L. Sustained hydrogen photoproduction by *Chlamydomonas reinhardtii*: effects of culture parameters. *Biotechnol. Bioeng.* **78**, 731–740 (2002).
- Kosourov, S., Patrusheva, E., Ghirardi, M. L., Seibert, M. & Tsygankov, A. A comparison of hydrogen photoproduction by sulfur-deprived *Chlamydomonas reinhardtii* under different growth conditions. *J. Biotechnol.* **128**, 776–787 (2007).

14. Kruse, O., Rupprecht, J., Mussgnug, J. H., Dismukes, G. C. & Hankamer, B. Photosynthesis: a blueprint for solar energy capture and biohydrogen production technologies. *Photochem. Photobiol. Sci.* **4**, 957–970 (2005).
15. Allahverdiyeva, Y. *et al.* Screening for biohydrogen production by cyanobacteria isolated from the Baltic Sea and Finnish lakes. *Int. J. Hydrogen Energy* **35**, 1117–1127 (2010).
16. Zehr, J. P. *et al.* Unicellular cyanobacteria fix N₂ in the subtropical North Pacific Ocean. *Nature* **412**, 635–638 (2001).
17. Welsh, E. A. *et al.* The genome of *Cyanothece* 51142, a unicellular diazotrophic cyanobacterium important in the marine nitrogen cycle. *Proc. Natl Acad. Sci. USA* **105**, 15094–15099 (2008).
18. Zehr, J. P., Methé, B. & Foster, R. in *Biological Nitrogen Fixation, Sustainable Agriculture and the Environment Vol. 41 Current Plant Science and Biotechnology in Agriculture* (eds Lin, M., Wang, Y.P., Tian, Z.X., Elmerich, C. & Newton, W. E.) 361–365 (Springer, 2005).
19. Schneegurt, M. A., Sherman, D. M. & Sherman, L. A. Growth, physiology and ultrastructure of the diazotrophic cyanobacterium, *Cyanothece* sp. strain ATCC 51142 in myxotrophic and chemotrophic cultures. *J. Phycol.* **33**, 632–642 (1997).
20. Stockel, J. *et al.* Global transcriptomic analysis of *Cyanothece* 51142 reveals robust diurnal oscillation of central metabolic processes. *Proc. Natl Acad. Sci. USA* **105**, 6156–6161 (2008).
21. Berman-Frank, I. *et al.* Segregation of nitrogen fixation and oxygenic photosynthesis in the marine cyanobacterium *Trichodesmium*. *Science* **294**, 1534–1537 (2001).
22. Toepel, J., Welsh, E., Summerfield, T. C., Pakrasi, H. B. & Sherman, L. A. Differential transcriptional analysis of the cyanobacterium *Cyanothece* sp. strain ATCC 51142 during light-dark and continuous-light growth. *J. Bacteriol.* **190**, 3904–3913 (2008).
23. Elvittigala, T., Stockel, J., Ghosh, B. K. & Pakrasi, H. B. Effect of continuous light on diurnal rhythms in *Cyanothece* sp. ATCC 51142. *BMC Genomics* **10**, 226 (2009).
24. Dutta, D., De, D., Chaudhuri, S. & Bhattacharya, S. K. Hydrogen production by cyanobacteria. *Microbial. Cell Factories* **4**, 36 (2005).
25. Feng, X. *et al.* Mixotrophic and photoheterotrophic metabolism in *Cyanothece* sp. ATCC 51142 under continuous light. *Microbiology* **156**, 2566–2574 (2010).
26. Eisbrenner, G. & Evans, H. J. Aspects of hydrogen metabolism in nitrogen-fixing legumes and other plant-microbe associations. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **34**, 105–136 (1983).
27. Liu, J. G., Bukatin, V. E. & Tsygankov, A. A. Light energy conversion into H₂ by *Anabaena variabilis* mutant PK84 dense cultures exposed to nitrogen limitations. *Int. J. Hydrogen Energy* **31**, 1591–1596 (2006).
28. Harwood, C. S. in *Bioenergy* (eds Harwood, C.S., Wall, J.D. & Demain, A.) 259–271 (ASM press, 2007).
29. Goldet, G. *et al.* Hydrogen production under aerobic conditions by membrane-bound hydrogenases from *Ralstonia* species. *J. Am. Chem. Soc.* **130**, 11106–11113 (2008).
30. Montoya, J. P. *et al.* High rates of N₂ fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean. *Nature* **430**, 1027–1032 (2004).
31. Mitsui, A. & Suda, S. Alternative and cyclic appearance of H₂ and O₂ photoproduction activities under nongrowing conditions in an aerobic nitrogen-fixing unicellular cyanobacterium *Synechococcus* sp. *Curr. Microbiol.* **30**, 1–6 (1995).
32. Schutz, K. *et al.* Cyanobacterial H₂ production—a comparative analysis. *Planta* **218**, 350–359 (2004).
33. Happe, T., Schutz, K. & Bohme, H. Transcriptional and mutational analysis of the uptake hydrogenase of the filamentous cyanobacterium *Anabaena variabilis* ATCC 29413. *J. Bacteriol.* **182**, 1624–1631 (2000).
34. Fay, P. Oxygen relations of nitrogen fixation in cyanobacteria. *Microbiol. Rev.* **56**, 340–373 (1992).
35. Houchins, J. P. & Burris, R. H. Comparative characterization of two distinct hydrogenases from *Anabaena* sp. strain 7120. *J. Bacteriol.* **146**, 215–221 (1981).
36. Tsygankov, A. A., Borodin, V. B., Rao, K. K. & Hall, D. O. H₂ photoproduction by batch culture of *Anabaena variabilis* ATCC 29413 and its mutant PK84 in a photobioreactor. *Biotechnol. Bioeng.* **64**, 709–715 (1999).
37. Tsygankov, A. A., Serebryakova, L. T., Rao, K. K. & Hall, D. O. Acetylene reduction and hydrogen photoproduction by wild-type and mutant strains of *Anabaena* at different CO₂ and O₂ concentrations. *FEMS Microbiol. Lett.* **167**, 13–17 (1998).
38. Borodin, V. B., Tsygankov, A. A., Rao, K. K. & Hall, D. O. Hydrogen production by *Anabaena variabilis* PK84 under simulated outdoor conditions. *Biotechnol. Bioeng.* **69**, 478–485 (2000).
39. Weyman, P. D., Pratte, B. & Thiel, T. Hydrogen production in nitrogenase mutants in *Anabaena variabilis*. *FEMS Microbiol. Lett.* **304**, 55–61 (2010).
40. Asada, Y. & Kawamura, S. Aerobic hydrogen accumulation by a nitrogen-fixing cyanobacterium, *Anabaena* sp. *Appl. Environ. Microbiol.* **51**, 1063–1066 (1986).
41. Xiankong, Z., Haskell, J. B., Tabita, F. R. & Baalen, C. V. Aerobic hydrogen production by the heterocystous cyanobacteria *Anabaena* spp. strains CA and 1F. *J. Bacteriol.* **156**, 1118–1122 (1983).
42. Shestakov, S. V. & Mikheeva, L. E. Genetic control of hydrogen metabolism in cyanobacteria. *Russian J. Genet* **42**, 1272–1284 (2006).
43. Kruse, O. *et al.* Improved photobiological H₂ production in engineered green algal cells. *J. Biol. Chem.* **280**, 34170–34177 (2005).
44. Min, H. T. & Sherman, L. A. Hydrogen production by the unicellular, diazotrophic cyanobacterium *Cyanothece* sp strain ATCC 51142 under conditions of continuous light. *Appl. Environ. Microbiol.* **76**, 4293–4301 (2010).
45. Reddy, K. J., Haskell, J. B., Sherman, D. M. & Sherman, L. A. Unicellular, aerobic nitrogen-fixing cyanobacteria of the genus *Cyanothece*. *J. Bacteriol.* **175**, 1284–1292 (1993).
46. Oda, Y. *et al.* Functional genomic analysis of three nitrogenase isozymes in the photosynthetic bacterium *Rhodospseudomonas palustris*. *J. Bacteriol.* **187**, 7784–7794 (2005).

Acknowledgments

We thank Lawrence Page for the initial standardization of the hydrogen measurements assay and all other members of the Pakrasi and Sherman laboratories for collegial discussions. We also thank Dr. Teresa Thiel for providing us with the wild-type *Anabaena variabilis* strain. This work was supported by funding from DOE-BER (DE-FC02-07ER64694) and from the Consortium for Clean Coal Utilization at Washington University.

Author contributions

A.B., J.S., L.A.S. and H.B.P. designed the experiments; A.B., J.S. and H.M. performed the experiments; A.B., J.S. and H.B.P. wrote the paper.

Additional information

Competing financial interests: The authors declare no competing financial interests.

Reprints and permission information is available online at <http://npg.nature.com/reprintsandpermissions/>

How to cite this article: Bandyopadhyay, A. *et al.* High rates of photobiological H₂ production by a cyanobacterium under aerobic conditions. *Nat. Commun.* 1:139 doi: 10.1038/ncomms1139 (2010).